



Analysis of cGMP signalling with transgenic mice expressing FRET-based cGMP sensors

Citation

Wen, L., M. Thunemann, S. Feil, M. Hillenbrand, A. Vachaviolos, T. Ott, X. Han, et al. 2013.
“Analysis of cGMP signalling with transgenic mice expressing FRET-based cGMP sensors.”
BMC Pharmacology & Toxicology 14 (Suppl 1): P76. doi:10.1186/2050-6511-14-S1-P76. <http://dx.doi.org/10.1186/2050-6511-14-S1-P76>.

Published Version

doi:10.1186/2050-6511-14-S1-P76

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:11877050>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

POSTER PRESENTATION

Open Access

Analysis of cGMP signalling with transgenic mice expressing FRET-based cGMP sensors

Lai Wen^{1†}, Martin Thunemann^{1†}, Susanne Feil¹, Matthias Hillenbrand¹, Angelos Vachaviolos¹, Thomas Ott², Xiaoxing Han³, Dai Fukumura³, Rakesh K Jain³, Cor de Wit⁴, Michael Russwurm⁵, Robert Feil^{1*}

From 6th International Conference on cGMP: Generators, Effectors and Therapeutic Implications
Erfurt, Germany. 28-30 June 2013

Background

Sensor proteins that detect cGMP are valuable tools to analyse the molecular mechanisms underlying cGMP's manifold physiological functions. These cGMP sensors allow for visualization and quantification of cGMP in living cells, in real time, and at high spatial and temporal resolution. For instance, they allow one to study sub-cellular cGMP compartments, and combined with multi-photon microscopy, they can provide new insights into complex biological processes, which can be analysed only in live animals.

Results

To monitor cGMP signals *in vivo*, we have generated and characterized transgenic mouse lines expressing the fluorescence resonance energy transfer (FRET)-based cGi500 sensor (cGMP indicator with an EC₅₀ value for cGMP of 500 nM, [1]). One mouse line produced by random transgenesis expresses cGi500 in vascular and visceral smooth muscle. Other mouse lines were created by targeted modification of the ROSA26 locus; these lines either show ubiquitous cGi500 expression, or they can be used for Cre/lox-dependent, tissue-specific cGi500 expression. These cGi500-transgenic mice are healthy and fertile, and do not show obvious adverse phenotypes.

Primary cells isolated from cGi500-expressing mice, including platelets, neural cells, and smooth muscle cells were used for FRET imaging experiments. NO-induced cGMP synthesis via soluble guanylyl cyclase was observed

in the majority of cell types, while cGMP responses to atrial or C-type natriuretic peptides differed between cell types. Moreover, NO-induced cGMP elevations could be observed in the vasculature of retinas isolated from cGi500-expressing animals. Importantly, the feasibility of FRET-based cGMP imaging in living animals was demonstrated by intravital microscopy of anesthetized cGi500-transgenic mice. By epifluorescence FRET microscopy we could visualize transient NO-induced cGMP signals in arterial walls of the cremaster muscle. Furthermore, we used multi-photon microscopy in a dorsal skinfold chamber model and were able to record NO-elicited FRET changes in subcutaneous vessels.

Conclusion

All in all, these cGMP sensor mouse lines represent valuable resources for studies with cGi500-expressing primary cells, and, importantly, they can also be used for intravital imaging studies with live animals. We believe that these mouse lines will find widespread applications and that in combination with other genetic mouse models they can contribute to improve our understanding of cGMP-modulated physiological or disease-associated processes.

Authors' details

¹Interfakultäres Institut für Biochemie, University of Tübingen, Tübingen, Germany. ²IZKF - Transgene Tiere, Universität Tübingen, Tübingen, Germany. ³Edwin L. Steele Laboratory for Tumor Biology, Department of Radiation Oncology, Massachusetts General Hospital (MGH) and Harvard Medical School, Boston, Massachusetts, USA. ⁴Institut für Physiologie, Universität zu Lübeck, Lübeck, Germany. ⁵Institut für Pharmakologie und Toxikologie, Universität Bochum, Bochum, Germany.

* Correspondence: robert.feil@uni-tuebingen.de

† Contributed equally

¹Interfakultäres Institut für Biochemie, University of Tübingen, Tübingen, Germany

Full list of author information is available at the end of the article

Published: 29 August 2013

Reference

1. Russwurm M, Mullershausen F, Friebe A, Jager R, Russwurm C, Koesling D: Design of fluorescence resonance energy transfer (FRET)-based cGMP indicators: a systematic approach. *Biochem J* 2007, **407**:69-77.

doi:10.1186/2050-6511-14-S1-P76

Cite this article as: Wen et al.: Analysis of cGMP signalling with transgenic mice expressing FRET-based cGMP sensors. *BMC Pharmacology and Toxicology* 2013 **14**(Suppl 1):P76.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

